

AD _____

GRANT NUMBER DAMD17-94-J-4449

TITLE: A Randomized Clinical Trial to Evaluate Advance Nursing
Care for Women with Newly Diagnosed Breast Cancer

PRINCIPAL INVESTIGATOR: Laurie Ritz, R.N.

CONTRACTING ORGANIZATION: Methodist Hospital and Park Nicollet
Medical Foundation
Minneapolis, Minnesota 55416

REPORT DATE: October 1996

TYPE OF REPORT: Annual

19970314 071

PREPARED FOR: Commander
U.S. Army Medical Research and Materiel Command
Fort Detrick, Frederick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for public release;
distribution unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

DUPLICATE INFORMATION

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE October 1996	3. REPORT TYPE AND DATES COVERED Annual (1 Oct 95 - 30 Sep 96)	
4. TITLE AND SUBTITLE A Randomized Clinical Trial to Evaluate Advance Nursing Care for Women with Newly Diagnosed Breast Cancer			5. FUNDING NUMBERS DAMD17-94-J-4449	
6. AUTHOR(S) Laurie Ritz, R.N.				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Methodist Hospital and Park Nicollet Medical Foundation Minneapolis, Minnesota 55416			8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Commander U.S. Army Medical Research and Materiel Command Fort Detrick, Frederick, Maryland 21702-5012			10. SPONSORING/MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution unlimited			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200) The purpose of this randomized clinical trial is to study the impact of an advanced practice nurse on the cost of care and quality of life for women newly diagnosed with breast cancer. The control group receives standard medical care while the experimental group receives standard medical care plus advanced nursing care. The number of participants who have been enrolled in this study is 146. Attrition (10%) has occurred at one half the projected rate and refusal to participate (29%) has been 11% lower than anticipated. Ineligibility due to a previous diagnosis of cancer and treatment at a different treatment site after diagnosis has been higher than anticipated. Modifications in the proposal have been submitted to increase our sample size. USAMRMC approval is pending. The cost model has been developed and costs identified within each of the categories. Data collection and entry are proceeding with a participant response rate of 94%. High data reliability and internal validity have been established.				
14. SUBJECT TERMS Breast Cancer , Advanced Practice Nurse Outcomes, Quality of Life, Cost-Effective Care			15. NUMBER OF PAGES 19	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT Unlimited	

FOREWORD

Opinions, interpretations, conclusions and recommendations are those of the author and are not necessarily endorsed by the U.S. Army.

LR Where copyrighted material is quoted, permission has been obtained to use such material.

Where material from documents designated for limited distribution is quoted, permission has been obtained to use the material.

Citations of commercial organizations and trade names in this report do not constitute an official Department of Army endorsement or approval of the products or services of these organizations.

In conducting research using animals, the investigator(s) adhered to the "Guide for the Care and Use of Laboratory Animals," prepared by the Committee on Care and use of Laboratory Animals of the Institute of Laboratory Resources, National Research Council (NIH Publication No. 86-23, Revised 1985).

LR For the protection of human subjects, the investigator(s) adhered to policies of applicable Federal Law 45 CFR 46.

In conducting research utilizing recombinant DNA technology, the investigator(s) adhered to current guidelines promulgated by the National Institutes of Health.

In the conduct of research utilizing recombinant DNA, the investigator(s) adhered to the NIH Guidelines for Research Involving Recombinant DNA Molecules.

In the conduct of research involving hazardous organisms, the investigator(s) adhered to the CDC-NIH Guide for Biosafety in Microbiological and Biomedical Laboratories.

Laurie J. Pitt 10/23/96
PI - Signature Date

TABLE OF CONTENTS

	<u>Page</u>
Front Cover	1
SF 298 Report Documentation Page	2
Foreword	3
Table of Contents	4
Introduction	5
Body	6-10
Conclusions	10
References	11-12
Appendix A	13
Appendix B	14-16
Appendix C	17-18
Appendix D	19

INTRODUCTION

NATURE OF PROBLEM AND BACKGROUND OF PREVIOUS WORK

Breast cancer accounts for almost one third of all cancers in women in the United States (US). Greater than 184,000 new cases will be diagnosed in 1996 alone. More than 44,000 women will die of breast cancer this year.¹ Cost-effective methods to manage breast cancer patients while continuing to achieve quality outcomes is a major US public health goal.

As costs decrease, it is unclear if quality outcomes are being maintained. In addition, factors including access to care, intricacy of the health care system, numerous caregivers, complexities of the diagnostic tests and procedures, and technical components of treatment can overwhelm patients and result in compromised quality outcomes.

An Advanced Practice Nurse (APN) could serve as a facilitator to ease the breast cancer patient's way through the health care system providing quality care in a cost-effective manner. The former Office of Technology Assessment (OTA) of the US Congress conducted a comprehensive review of 286 studies on the cost and effectiveness of APNs. Their findings from this review indicated that within the APN's area of competence, they communicate better with patients, concentrate more on prevention, and provide more education than physicians. Patients are satisfied with care, access to care is less complicated, and the costs of care are less with the interventions of the APN.²

Additional studies including those which focused on lung cancer patients, low birthweight infants, MI patients, HIV-infected individuals, children with chronic diseases, and hospitalized elderly, have demonstrated the effectiveness of advanced nursing care with results of improved outcomes and reduced health care costs, but no one has focused on women with breast cancer.³⁻¹⁰

PURPOSE OF PRESENT WORK AND METHOD OF APPROACH

The purpose of this study is to test the following hypotheses:

- Women with newly diagnosed breast cancer who receive advanced nursing follow-up care/interventions will have a lower cost of care than patients who do not receive advanced nursing care.
- Women with newly diagnosed breast cancer who receive continuity of care through advanced nursing care/interventions across the various health care settings will achieve a better quality of life than patients who do not receive advanced nursing care.

The method of approach for this study is a randomized clinical trial to test these hypotheses using Brooten's model,¹¹ with modifications, to provide the framework for studying cost and quality of patient outcomes with the interventions of APNs.

BODY

Previously Reported Work

Preparation to Begin Study

As defined in the statement of work, hiring of staff, determination of APN charges, preparation of the data collection tools, and initial as well as continuing education of all participating staff has occurred as previously reported in the annual report for the period of October, 1994 - September, 1995.

Cost Analysis Model

The Cost Analysis Model has been completed (Appendix A). It consists of two main branches (total charges and total reimbursement) which continually divide to take into account the desired outcomes of the analysis of the study such as in-system versus out-of-system costs, and cancer related versus non-cancer related costs. The final extension of this model creates the cost categories of hospitalizations, emergency room visits, outpatient services, and non-charge estimates. The costs included within each of these final categories has also been identified. (Appendix B).

Accrual, Intervention, and Data Collection

Women with a newly diagnosed breast cancer are identified through positive pathology reports and referral by their physicians. They are then approached about their participation in the study, eligibility criteria are reviewed, informed consent is obtained and participants are randomized to one of two groups. Women in the control group receive the standard care which they would receive in the clinic, hospital, or patient's home. Women enrolled in the experimental group have an APN who begins follow-up immediately and continues follow-up for two years. All participants are asked to complete quality of life questionnaires at intervals of one week, one, three, six, twelve, eighteen, and twenty four months as well as a patient diary (schedule of their episodes of care) for the two year period.

Current Work (October 1995-September 1996)

Accrual Status

Accrual status to date is summarized in Table 1 as follows:

OCTOBER 1, 1994 - SEPTEMBER 30, 1996	Number of patients
Newly diagnosed breast cancer patients	419
Referred	361 (86%)
Eligible	206 (57%)
Refused	60 (29%)
Participants enrolled from eligible	146 (71%)
Lost to attrition	15 (10%)
Total participants	131

Four hundred nineteen women were diagnosed with breast cancer within our system of care during the first two years of study (October 1, 1994 - September 30, 1996). One hundred fifty-five were ineligible with sixty refusing to participate. The refusal rate (29%) is lower than anticipated, when compared with Hughe's reported refusal rate of 40% at the time of diagnosis.¹² Attrition (10%) has occurred at a rate of one half the projected rate.

Ineligibility is higher than anticipated. Reasons for study ineligibility during the past year include: previous diagnosis of cancer (n= 37), going outside the approved site of the study for care after diagnosis (n=28), inability to enroll within two weeks after diagnosis (n= 12), comorbidity which limits functional ability (n=7), inability to give informed consent (n=3), inability to complete questionnaires (n=3), severe psychiatric illness (n= 2), and inability to read or write English (n=1).

A modification to the study which increased the registration time of the subjects from one to two weeks following diagnosis was approved 10/16/95. This modification has increased our rate of enrollment from 6.8 participants/month to 7.3 participants /month.

Changes in health care delivery are occurring within our system. The diagnosis of breast cancer is made at our site with an increasing number of women choosing another hospital for treatment. Another change is the addition of 70,000 patients to our system of care beginning January 1, 1997. Based on these changes, we have written and submitted two further modifications to this study.

The first modification would allow enrollment of women who are being treated at a second hospital site. The second modification would extend our study period from 9/30/98 to 9/30/99. These modifications will increase our sample to 100 subjects in each group which using an alpha of .05 and statistical power of .8, will produce statistically significant results.

Although we are requesting extension of the study period, we are not asking for additional funding. Approval of the additional study site and extension of our period of study has been approved by the IRB both at our current and proposed study sites and by our grant administrator. Approval from the USAMRMC is pending.

APN Intervention

Follow-up care and interventions of the APN are based on Brooten's work¹¹ and the standards of advanced practice in oncology nursing.¹³ It includes coordination of care, assessment and monitoring of symptoms, direct care, patient and family education, consultation with other health care services, utilization of current research findings, and establishment of standards of practice. Care is individualized to the patient and family needs, based on the expressed needs of the individual, the assessment of the APN, and other health care providers' evaluations. A detailed description of the standard APN follow-up care is listed in Appendix C.

Data Collection and Entry

Collection of data is proceeding as outlined in the protocol. The response rate for the sets of the questionnaires (week 1, month 1, month 3, month 6, and monthly 12) for participants currently enrolled on the study is 94% (431/460). The data from the questionnaires has been continually entered into their appropriate databases as the sets are returned.

Collection methods were set up to gather the specific cost information from the hospital's billing system (Phamis), the clinic's billing system (Med.I.C.), and independent physicians affiliated with the hospital. Phamis is a health information management system and serves as the patient information data repository for the hospital. It runs on a Tandem midrange computer and consists of a variety of modules including encounter/visit management, ancillary management, nursing management, medical record management, patient registration, laboratory results reporting, pharmacy management, and patient accounting. Med.I.C. is a clinical management application that runs on an IBM 3090 developed by the Management Information System's (MIS) staff at Park Nicollet Clinic. It assumes the same role as Phamis in the clinical setting.

The collection methods for Phamis and Med.I.C. involve querying each system by the patient's chart number and outputting the desired data into a file with a delimited text format. This file is then imported into the correct database. The independent physicians are providing their cost information through their billing systems and this data is entered into a database. In order to carry out the planned analyses the detailed cost data must then be formed into a composite database summarizing in-system, out-of-system, cancer-related, and non-cancer-related costs for each patient in order to carry out the planned analyses. A mock cost dataset was used to create and test an algorithm (SAS) for reducing the multiple records to an analyzable dataset containing only 8 records per person. These 8 records include charge and reimbursement amounts for each of the four categories listed above.

Non-charge estimates are collected from APN logs and patient diary data. The data collected in the APN logs will be applied to a standardized APN intervention estimate. This APN intervention estimate is determined by using mean salary and benefits to calculate a per minute cost for time spent with each patient in the intervention arm. The other estimates will not have charges assigned to them and will be reported separately.

The collection methods of treatment and tumor information are also established. This information is abstracted from the hospital's oncology registry. The registry is a data system designed for the collection, management and analysis of data on persons with the diagnosis of cancer.¹⁴ The data is abstracted through a query of the registry's database and exported as a delimited file. This file is then imported into the correct database.

Databases have been developed and currently exist for the entry of questionnaire information, diary information, APN log data, treatment and tumor information, and the cost information from participating physicians, clinics, and hospital(s). These databases are related to one another via a variety of relational variables. These relational variables are unique identifiers for each patient in Phamis, Med.I.C., and the oncology registry and are indexed to the patient's study identification (ID) in a database designed specifically for that purpose. Data dictionaries have also been created to identify the variables in each of these databases.

The analysis of the data will begin in month 48 of the current statement of work or month 56 of the proposed work agreement.

Data Quality

The reliability of the data collected from the patient diary was evaluated to determine areas of strength and weakness. Each ledger-format diary allows patients to record data about episodes of care and participant/family members days lost from work and child care. Responses (n=18) from the first patient diaries (0-6 months) were compared to the hospital-clinic computer system to determine validity. The mean perfect agreement was

9.06 (standard error (steer) = 0.366) on a binary score of 11 items. The intervention group (score 9.09, stderr=0.495) had a greater number of patients with perfect agreement (correct physician/nurse name and correct date) for the categories including: primary doctor visits (8/11 vs. 4/7), surgery visits (4/11 vs. 2/7), plastic surgery visits (11/11 vs. 6/7), other visits (7/11 vs. 2/7), and urgent care visits (11/11 vs. 6/7). The control group (score 9.0 stderr 0.577) had a greater number of patients with perfect agreement for the hospitalizations (7/7 vs. 10/11), medical oncologist visits (4/7 vs. 5/11), nursing visits (6/7 vs. 8/11), nurse practitioner (6/7 vs. 9/11), home care visits (7/7 vs. 10/11), emergency room visits (7/7 vs. 10/11) and treatment category (6/7 vs. 7/11). Individual visits to health professionals and the overall treatment categorization show the most discordance. The patient diaries report fewer visits compared to the hospital-clinic computer system. Our long-term diary is a useful data collection tool (less than 10% disagreement) for breast cancer patients for homecare visits, hospitalizations, urgent care or emergency room visits. It can provide data about support group attendance, out-of-system hospital and clinic visits, child care use and days of work lost. This information is unavailable through other common data sources and is necessary to conduct the planned cost-benefit analysis.

CONCLUSIONS

Women with a newly diagnosed breast cancer who meet the study criteria, are participating in the study at a higher than initially projected rate. The refusal rate is 11% less than projected and attrition is one half of the projected rate. The questionnaire response rate is 94%. Comparisons of data from multiple sources indicate high reliability and internal validity.

The Cost Analysis Model is completed and is providing the framework to collect patient charges, reimbursement, and nonbillable costs. The APN intervention is being successfully implemented. Standard follow-up is well-defined with additional cares based on the patient's needs. Results of the completed data have not been analyzed but many anecdotal responses of the participating women, their health care providers, and other patients not participating in this study indicate positive outcomes.


Ineligibility of women has contributed to a lower than expected rate of enrollment, but protocol changes have been recommended to address this issue. These changes are feasible and will not alter the scientific merit of the study. With the USAMRMC's approval of the addition of another study site and another year of study (no additional funding requested), we will meet the work agreement as defined in our statement of work.

REFERENCES

1. Parker, SL; Tong, T.; Bolden, S; & Wingo, PA. (1996). Cancer statistics, 1996. CA: A Cancer Journal for Clinicians, 46(1), 5-27.
2. Office of Technology Assessment, U.S. Congress. (1986). Nurse Practitioners, Physician Assistants, and Certified Nurse-Midwives: A Policy Analysis. (Health Technology Case Study 37), OTA-HCS-37. Washington, DC: U.S. Government Printing Office.
3. Oleske, DM & Hauck, WW (1988)). A population-based evaluation of the impact of interventions for improving care to cancer patients in home settings. Home Health Services Quarterly. 9, 45-61.
4. McCorkle, R; Benoliel, JQ; Donaldson, G; Georgiadou, F; Moinpour, C; & Goodell, B. (1989). A randomized clinical trial of home nursing care of lung cancer patients. Cancer, 64, 1375-1382.
5. Brooten, D; Kumar, S; Brown, LP; Butts, P; Finkler, SA; Bakewell-Sachs, S; Gibbons, A; & Delivoria-Papadopoulos, M. (1986). A randomized clinical trial of early hospital discharge and home follow-up of very low-birth-weight infants. The New England Journal of Medicine, 315, 934-939.
6. Pozen, MW; Stechmiller, JA; Harris, W; Smith, S; Fried, DD; & Voigt, GC. (1977). A nurse rehabilitator's impact on patients with myocardial infarction. Medical Care, 15(10), 830-837.
7. Aiken, LH; Lake, ET; Semann, S; Lehamn, HP; O'Hare, PA; cole, CS; Dunbar, D; & Frank, I. Nurse practitioner managed care for persons with HIV infection. Image: Journal of Nursing Scholarship, 25(3), 172-177.
8. Lipman, T. (1986). Length of hospitalization of children with diabetes: Effect of a clinical nurse specialist. The Diabetes Educator, 14, 41-43.
9. Alexander, JA; Younger, RE; Cohen, RM; & Crawford, LV. (1988). Effectiveness of a nurse managed program for children with chronic asthma. Journal of Pediatric Nursing, 3, 312-317.
10. Neidlinger, L; Scroggins, K; & Kennedy, L. (1987). Cost evaluation of discharge planning for hospitalized elderly. Nursing Economics, 5, 225-230.

11. Brooten, D; Brown, LP; Munro, BH; York, R; Cohen, SM; Roncoli, M; & Hollingworth, A. (1988). Early discharge and specialist transitional care. Image: Journal of Nursing Scholarship, 20(2), 64-68.
12. Hughes, KK. (1993). Decision making by patients with breast cancer: The role of information in treatment selection. Oncology Nursing Forum, 20(4), 623-628.
13. Oncology Nursing Society. (1990). Standards of Advanced Practice in Oncology Nursing, Oncology Nursing Press, Inc.
14. National Cancer Registrars Association, Inc. Fact Sheet: The cancer registry and the registrar. Mundelein, IL: National Office.

Cost Model



Cost Model

Identification of Costs Included Within Each Category

1. Hospitalizations

A) Inpatient

- a) Provider procedures and services
- b) Room utilization
- c) Radiologic procedures
- d) Laboratory tests
- e) Supplies
- f) Medications; Drugs
- g) Rehab services (if needed)
- h) ER services (if needed)
- i) Professional fees

Nurse anesthetist
EKG reading by cardiologist
Park Nicollet physician fees

B) 23 hr

same as inpatient charges

C) One day surgery

same as inpatient charges

2. Emergency room visit charges

- A) Provider procedures and services
 - B) Radiologic procedures
 - C) Laboratory tests
 - D) Supplies
 - E) Medications; Drugs
 - F) ER professional fees
- Park Nicollet physician fees

3. Outpatient service charges

- A) Park Nicollet clinic visit charges
 - a) Provider procedures and services
 - b) Radiologic procedures
 - c) Laboratory tests
 - d) Supplies
 - e) Medications; Drugs
- B) Outpatient hospital service charges
 - a) Chemotherapy visit charges
 - i) Provider procedures and services
 - ii) Radiologic procedures
 - iii) Laboratory tests
 - iv) Supplies
 - v) Medications; Drugs
 - b) Radiation therapy visit charges
 - same as chemotherapy charges
 - Radiation oncologist fees
 - c) Home care visit charges
 - same as chemotherapy charges
 - d) Rehabilitation visit charges
 - same as chemotherapy charges

4. Non-charge estimates

- A) Advanced Practice Nurse intervention
 - a) Patient visits
 - b) Home care visits
 - c) Telephone calls
 - d) Administrative
 - e) Travel

B) Rehabilitation services

- a) Breast cancer support groups
- b) Other support groups
- c) Chaplain
- d) Psychologist (out-of-system)
- e) Social worker
- f) Sex therapist (out-of-system)
- g) Dietitian

C) Telephone calls

- a) Nurse
- b) Physician

- D) Child care / lost work time
- E) Hospitalizations (out-of-system)
- F) Home Care visits (out-of-system)
- G) Referring physicians (out-of-system)

Standard APN Follow-up Care

PHASE I INTRODUCTION		PHASE III FOLLOW-UP IF NO TREATMENT	
Frequency	Pre-surgical meeting @ 0-7 days	Frequency	Radiation, Chemo, Surgery, Plastic Surgery
	Introductory meeting		Weekly contact for status
1visit; 1 call	Explanation of BCNC role & availability	weekly	Education, support and assessment
1visit; 1 call	Needs assessment form	x 2-24wks	Pain
1visit; 1 call	Decision making process	x 2-24wks	ROM
1visit; 1 call	Physical assessment form with Hx (PRN)	x 2-12wks	Seroma
hospital visit	Give pt. copy history/current meds	x 2-12wks	Necrosis
hospital visit	Library information given	x 2-26 wks	Oral intake (especially with chemo)
hospital visit		x 2-12wks	Infection
	Follow-up-up plan:	ongoing	Fatigue
	Tentative plan of care:	x 2-26 wks	Prosthesis Information
	Obtain arm measurements bilaterally	x 2-26 wks	Blood counts
daily callx1-2wks	Calendar	x 2-26 wks	Psychosocial support
wkly callsx6-8	Accompany to MD visits		Mood
1 visit	Next contact with BCNC (date)	ongoing	Coping
ongoing	Contact during hospitalization	ongoing	Energy level
ongoing	Contact during outpatient visit	ongoing	Referral to Social Services PRN
			Referral to Support Groups in community
1-2 visits	PHASE II POST OP	ongoing	BCNC support during any/all visits
1-3x/week	Home visit post-op 24-48 hrs	ongoing	surgeon, plastics, oncologist, radiation
	Telephone contact during 1st 3-5 days		
	Education	ongoing	Follow-up visit @ 4-6 weeks (all pts)
x 1	Signs of infection/inflammation		Physical assessment
daily x 7	Temp		Arm measurements
daily x 7	JP Stripping /Drainage / leakage/	x1-2 visits	Review signs/symptoms of lymphedema
daily x 7	Incisional Pain	PRN	Body image-looked in mirror?
daily x 7	Swelling	x1 or PRN	Prosthesis
daily x 7	Redness	x1 or PRN	Sexuality
daily x 7	Arm ROM/ pain / burning	x1 or PRN	Back to work or normal activity yet?
daily x 7	General well-being	x1 or PRN	Told others?
daily x 7	Mood	1-24 wks	Family
daily x 7	Fatigue	1-4 wks	Friends
daily x 7	Energy level	1-4 wks	Co-workers
daily x 7	Appetite	1-4 wks	Support group?
daily x 7	Comfort/pain control/ constipation	1-24 wks	Follow-up with oncologist
weekly	Coping with life and home	ongoing	Support
weekly	Spouse/significant other	ongoing	Treatment discussion
weekly	Family issues	ongoing	Options
weekly	Children	ongoing	Reinforce education
weekly	Child care	ongoing	Wigs
weekly	Job/career	2-24wks	Cosmetics/hair care
weekly	Housework	ongoing	Fatigue management
X1	Have they met with reach to recovery	ongoing	Hot flashes and management
weekly	Exercise/ review with pt /	ongoing	Follow-up with Plastic Surgeon
x2	Prosthesis	ongoing	Monitor for necrosis
	Follow-up surgeon visit date?	ongoing	Monitor for infection
5 visits/ ongoing	Medical plan of care:	ongoing	Assess for normal ADL's
	RT	ongoing	Pain control with saline expansion
	Chemo	ongoing	Plan for secondary surgery PRN
	Additional Surgery		
	Next FU visit scheduled?		

Standard APN Follow-up Care

PHASE III TREATMENT MANAGEMENT		PHASE IV FOLLOW-UP CARE	
Frequency		Frequency	
monthly FU	Tamoxifen		Telephone contact every other wk x 4
0-2 yrs	Side effects: Hot flashes, weight, mood swings, endometrial ca risk	ongoing	(every week x 4 if no treatment; then qow
	GYN evaluation if spotting	ongoing	Monthly FU phone calls or visits for all pts
daily x1-3	Chemo: Call day 1,2,3	ongoing	Lymphedema FU every 3 mos x 4; then q 6
weekly 0-32	Assess nausea, fatigue, diet, activity,		BSE instruction with return demo PRN
& monthly	diarrhea, constipation, mouth sores	ongoing	give shower cards, stickers
	Blood counts	ongoing	Mammogram scheduled annually
	Educate regarding plan & ttment delays	monthly	Stress importance of BSE and FU care
FU weekly	Radiation Therapy:	ongoing	ISSUES: support, assess and educate
0-10 weeks	Assess skin reaction, fatigue,	ongoing	Diet
	blood counts	ongoing	Exercise
0-10 weeks	Educate regarding ttment plan and FU	ongoing	Weight
		ongoing	Hot flashes
monthly	Educational reinforcement	ongoing	Sexuality
ongoing	Frequency of healthcare visits:	ongoing	Pregnancy
ongoing	Strategy for coping	ongoing	Work Issues
ongoing	Activity adjustment	ongoing	Menopause
ongoing	Fatigue management	ongoing	Insurance coverage
		ongoing	Medication cost
monthly	Activity of daily life	ongoing	Venous access device management
ongoing	Ability to perform ADL's	ongoing	Late treatment effects
ongoing	Appearance	monthly	Health Promotion
ongoing	Fatigue	ongoing	Quit smoking
ongoing	Energy level	ongoing	Diabetic control
ongoing	Change from precancer level of activity	ongoing	Assess hypertension
ongoing	Diet adjustment	ongoing	Dietary modifications
ongoing	Oral rinse and mouth care	ongoing	Stress reduction
ongoing	Fluid intake	monthly	Complementary therapies
ongoing	Monitor output	ongoing	Stress management
ongoing	Taste changes	ongoing	Imagery
ongoing	Weight gain/loss	ongoing	Positive thinking
ongoing	Social adjustment	ongoing	Support groups
ongoing	Sick leave availability	monthly	Recovery
ongoing	Child care issues	ongoing	Taking control/proactive
ongoing	Transportation to treatment	ongoing	Fear of recurrence
ongoing	Cooking	ongoing	Coping
ongoing	Cleaning	ongoing	Spirituality
ongoing	Laundry	ongoing	Hope
ongoing	Shopping	monthly	Psychosocial assessment:
ongoing	other	ongoing	Kids
monthly	Physical side effects	ongoing	Sex
ongoing	Skin care:	ongoing	Work
ongoing	Rashes	ongoing	Home
ongoing	Incision	ongoing	Reconstruction
ongoing	Dryness	ongoing	Future plans
ongoing	Neuropathy	ongoing	Social Services referral PRN
ongoing	Status of surgical site		

ACRONYM AND ABBREVIATION DEFINITIONS

APN	-	Advanced Practice Nurse
EKG	-	Electrocardiogram
ER	-	Emergency Room
HIV	-	Human Immunodeficiency Virus
MI	-	Myocardial Infarction
MIS	-	Management Information Systems
OTA	-	Office of Technology Assessment